

Managed Long Term Care Quality Indicator Benchmarks, July - December 2016 Data*												
			Percentile					Incentive year				
Type	Measure	Statewide Rate	0	50	75	90	100	2014	2015	2016	2017	2018
Prevalence	Risk-adjusted percentage of members who did not have an emergency room visit in the last 90 days [†]	92.1	79.8	91.0	93.2	95.1	96.9	✓	✓	✓	✓	
Prevalence	Risk-adjusted percentage of members who did not have falls that required medical intervention in the last 90 days [†]	94.2	78.4	94.0	95.2	95.9	98.2	✓	✓	✓	✓	
Prevalence	Percentage of members who managed their medications independently [†]	10.2	0.3	10.7	16.3	27.0	31.9					
Prevalence	Risk-adjusted percentage of members who did not experience uncontrolled pain	88.0	64.9	85.7	92.0	95.7	97.6		✓	✓	✓	
Prevalence	Risk-adjusted percentage of members who did not experience severe or more intense pain daily	95.2	68.2	94.7	97.7	98.4	98.8	✓				
Prevalence	Risk-adjusted percentage of members who were not lonely or were not distressed	91.9	67.9	90.9	93.9	94.7	98.7	✓	✓	✓	✓	
Prevalence	Percentage of members who received an influenza vaccination in the last year [†]	76.9	69.4	78.0	81.6	87.2	94.4	✓	✓	✓	✓	
Prevalence	Percentage of members who responded that a health plan representative talked to them about appointing someone to make decisions about their health if they are unable to do so [†]	64.7	41.7	70.2	76.7	79.0	92.9		✓	✓	✓	
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in ADL function	84.7	72.8	85.8	89.2	93.2	100.0					
Over-Time	Percentage of members who remained stable or demonstrated improvement in IADL function	87.8	69.5	86.1	89.8	92.6	95.4					
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in cognition	76.7	58.8	79.1	81.8	86.0	93.8					
Over-Time	Percentage of members who remained stable or demonstrated improvement in communication	82.9	71.1	84.4	87.1	89.0	93.8					
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in pain intensity	86.6	66.3	85.8	88.0	90.9	94.6		✓	✓	✓	
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in mood	83.1	57.1	82.9	85.4	88.2	94.2					
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in NFLOC score [†]	85.8	68.2	86.6	88.8	91.9	95.7		✓	✓	✓	
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in locomotion [†]	68.0	58.1	70.7	81.7	92.1	99.3					
Over-Time	Percentage of members who remained stable or demonstrated improvement in bathing [†]	73.7	58.7	73.2	77.6	79.5	90.4					
Over-Time	Percentage of members who remained stable or demonstrated improvement in toilet transfer [†]	69.6	49.6	71.2	76.6	81.1	84.8					
Over-Time	Percentage of members who remained stable or demonstrated improvement in dressing upper body [†]	75.2	55.8	74.5	77.5	81.1	89.1					
Over-Time	Percentage of members who remained stable or demonstrated improvement in dressing lower body [†]	71.7	52.3	70.8	75.9	79.9	87.5					
Over-Time	Percentage of members who remained stable or demonstrated improvement in toilet use [†]	64.9	42.7	68.4	73.0	78.2	90.4					
Over-Time	Percentage of members who remained stable or demonstrated improvement in eating [†]	79.6	62.3	81.2	84.5	88.8	93.6					
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in urinary continence [†]	74.8	61.1	75.7	78.0	80.1	82.2		✓	✓	✓	
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in shortness of breath [†]	88.9	80.0	86.8	90.5	93.2	96.2		✓	✓	✓	
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in managing medications [†]	66.7	44.4	65.6	69.2	75.2	81.0					
*Excluding nursing home residents and first assessments (assessment reason= first assessment)												
† New York created.												