

Managed Long Term Care Quality Indicator Benchmarks, July - December 2018 Data*													
			Percentile					Incentive year					
Type	Measure	Statewide Rate	0	50	75	90	100	2014	2015	2016	2017	2018	2019
Prevalence	Risk-adjusted percentage of members who did not have an emergency room visit in the last 90 days	93.4	80.9	92.6	94.4	95.0	96.6	✓	✓	✓	✓	X	✓
Prevalence	Risk-adjusted percentage of members who did not experience falls that resulted in major or minor injury in the last 90 days	95.3	84.4	95.3	95.7	96.4	100.0					✓	✓
Prevalence	Risk-adjusted percentage of members who did not have falls that required medical intervention in the last 90 days	--	--	--	--	--	--	✓	✓	✓	✓		
Prevalence	Percentage of members who managed their medications independently	5.7	0.2	7.9	13.6	24.4	31.4						
Prevalence	Risk-adjusted percentage of members who did not experience uncontrolled pain	96.3	77.5	95.9	98.2	99.6	100.0		✓	✓	✓	✓	✓
Prevalence	Risk-adjusted percentage of members who did not experience severe or more intense pain daily	98.6	81.1	99.1	99.7	99.9	100.0	✓					
Prevalence	Risk-adjusted percentage of members who were not lonely or were not distressed	97.2	74.5	96.9	99.1	99.7	100.0	✓	✓	✓	✓	✓	✓
Prevalence	Percentage of members who received an influenza vaccination in the last year	78.7	68.3	79.8	85.0	87.5	93.0	✓	✓	✓	✓	✓	✓
Prevalence	Percentage of members who responded that a health plan representative talked to them about appointing someone to make decisions about their health if they are unable to do so	74.4	59.4	77.1	78.6	86.3	97.0		✓	✓	✓	✓	✓
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in ADL function	88.5	74.2	88.4	90.5	92.0	94.3						
Over-Time	Percentage of members who remained stable or demonstrated improvement in IADL function	91.0	71.4	89.7	93.6	95.0	97.8						
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in cognition	80.1	60.5	79.7	84.8	87.6	91.3						
Over-Time	Percentage of members who remained stable or demonstrated improvement in communication	83.7	70.2	84.4	86.8	88.2	96.1						
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in pain intensity	90.7	79.8	90.0	92.7	93.6	96.9		✓	✓	✓	✓	✓
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in mood	85.5	69.3	85.5	87.0	88.8	93.5						
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in NFLOC score	89.5	70.5	89.5	91.6	94.3	95.5		✓	✓	✓	✓	✓
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in locomotion	74.2	48.2	74.6	84.0	94.6	100.0						
Over-Time	Percentage of members who remained stable or demonstrated improvement in bathing	79.1	59.0	76.8	80.8	85.6	89.6						
Over-Time	Percentage of members who remained stable or demonstrated improvement in toilet transfer	76.1	50.0	75.5	79.0	84.9	92.1						
Over-Time	Percentage of members who remained stable or demonstrated improvement in dressing upper body	80.0	65.9	77.0	80.2	85.8	89.2						
Over-Time	Percentage of members who remained stable or demonstrated improvement in dressing lower body	78.4	53.6	75.0	78.7	83.8	89.2						
Over-Time	Percentage of members who remained stable or demonstrated improvement in toilet use	72.3	39.3	72.0	76.2	84.5	90.7						
Over-Time	Percentage of members who remained stable or demonstrated improvement in eating	80.2	61.8	81.6	85.0	90.4	92.9						
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in urinary continence	82.3	72.6	81.3	85.1	87.3	91.0		✓	✓	✓	✓	✓
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in shortness of breath	92.6	80.9	92.9	95.2	96.0	98.0		✓	✓	✓	✓	✓
Over-Time	Risk-adjusted percentage of members who remained stable or demonstrated improvement in managing medications	71.6	46.7	71.7	74.3	80.5	82.5						
*Excluding nursing home residents and first assessments (assessment reason= first assessment)													